

Record of flocks band of Black-bellied whistling duck, *Dendrocygna autumnalis* (Linnaeus, 1758) (Anseriformes: Anatidae) with *Dendrocygna viduata* (Linnaeus, 1766) and *Dendrocygna bicolor* (Vieillot, 1816)

Registro de formação de bando misto de marreca-cabocla, *Dendrocygna autumnalis* (Linnaeus, 1758) (Anseriformes: Anatidae) com seus conge-néricos *Dendrocygna viduata* (Linnaeus, 1766) e *Dendrocygna bicolor* (Vieillot, 1816)

Aloysio Souza de Moura^{1,2*}, Felipe Santana Machado¹, Ravi Fernandes Mariano¹, Henrique Siqueira¹, Bruno Senna Corrêa² & Marco Aurélio Leite Fontes¹

1 Departamento de Ciências Florestais (DCF), Laboratório de Ecologia Florestal. Caixa postal 197, CEP 37.200-000, Lavras, MG. 2 CEFET Nepomuceno, Av. Monsenhor Luiz de Gonzaga, 103, Centro, CEP 37250-000, Nepomuceno/MG.

*Corresponding author: thraupidaelo@yahoo.com.br

Resumo Sick menciona em sua obra “Ornitologia Brasileira” que os anatídeos são aves sociáveis que podem ocorrer em grandes concentrações, porém, menciona que bandos da espécie *Dendrocygna autumnalis* não se mesclam com irerês (*Dendrocygna viduata*) e outras marrecas. Apresentamos aqui este registro atípico e ainda desconhecido na literatura científica.

Palavras-chave: Anatidae, bandos mistos, registro

Abstract Sick mentions in your book “Ornitologia Brasileira” that the anatids are sociable birds with high number of individuals in its populations. Furthermore, also mentions that *Dendrocygna autumnalis* bird flock do not mix with white-faced whistling duck (*Dendrocygna viduata*) and others ducks. we

present this atypical record still unknown in scientific literature.

Keywords: Anatidae, mixed flocks, record

Between the 26 bird species in Anatidae from Brazil (PIACENTINI et al. 2015), it is allocated the taxon popularly known as black-bellied whistling duck, *Dendrocygna autumnalis* (Linnaeus, 1758). Sick (1997) mentions in your book “Ornitologia Brasileira” that the anatids are sociable birds with high number of individuals in its populations. Futher more, also mentions that *D. autumnalis* bird flock do not mix with white-faced whistling duck (*Dendrocygna viduata*) and others ducks.

We aim to contribute with the knowledge of natural history, we present here a mixed band com-

posed by *Dendrocygna autumnalis*, *Dendrocygna viduata* and *Dendrocygna bicolor* (Vieillot, 1816). During the March 29th, 2008, around 03h15 PM, we recorded and photographed a mixed flocks of anatids composed by species mentioned above, in Jequitaí city (Figure 1), north of Minas Gerais State, southeastern Brazil (Figure 2).

The birds are foraging in a natural lake near to the edge of Jequitaí river ($17^{\circ}11'36.57''S/44^{\circ}28'45.66''O$, 495m), and when detected our presence, the birds flying together in a great mixed flocks.

With the objective to contribute of the knowledge of ecology and habits of the specie, we present this atypical record still unknown in scientific literature.




Figura 1. Location of the municipality of Jequitaí, north of the state of Minas Gerais, south east Brazil.




Figura 2. Mixed flocks composed by *Dendrocygna viduata* (left), *Dendrocygna bicolor* (center) and *Dendrocygna autumnalis* (right), Jequitaí city, north of Minas Gerais State, Southeastern Brazil. (Photo by Aloysio S. de Moura).

References

Piacentini VQ, Aleixo A, Agne CE, Maurício GN, Pacheco JF, Bravo GA, Brito GRR, Naka LN, Olmos F, Posso S, Silveira LF, Betini GS, Carrano E, Frans I, Lees AC, Lima LM, Pioli D, Schunk F, Amaral FR, Bencke GA, Cohn-Haft M, Figueiredo LFA, Straube FC, Cesari E (2015) Annotated checklis to fthe birds of Brazil by the Brazilian Ornithological Records Committee / Lista comentada das aves do Brasil pelo Comitê Brasileiro de Registros Ornitológicos. **Revisão Brasileira de Ornitologia** 23(2): 91-298.

SICK, H. (1997) **Ornitologia Brasileira**. Rio de Janeiro: Editora Nova Fronteira. 862p.